

Use of pyrophyllite to reduce heavy metals mobility in a soil environment

S. Murtić^{1,*}, H. Čivić², E. Sijahović², I. Koleška³, J. Jurković⁴ and M. Tvica⁵

¹University of Sarajevo, Faculty of Agriculture and Food Sciences, Department of Plant Physiology, Zmaja od Bosne 8, BA71000 Sarajevo, Bosnia and Herzegovina

²University of Sarajevo, Faculty of Agriculture and Food Sciences, Department of Plant Nutrition, Zmaja od Bosne 8, BA71000 Sarajevo, Bosnia and Herzegovina

³University of Banja Luka, Faculty of Agriculture, Department of Plant Physiology and Nutrition, Bulevar vojvode Petra Bojovića 1A, BA78000 Banja Luka, Bosnia and Herzegovina

⁴University of Sarajevo, Faculty of Agriculture and Food Sciences, Department of Chemistry, Zmaja od Bosne 8, BA71000 Sarajevo, Bosnia and Herzegovina

⁵University of Sarajevo, Faculty of Agriculture and Food Sciences, Department of Pedology, Zmaja od Bosne 8, BA71000 Sarajevo, Bosnia and Herzegovina

*Correspondence: murticsenad@hotmail.com

Abstract. This study revealed the effects of pyrophyllite ore materials on heavy metals mobility in soil plots located near the steel mill in Zenica (Bosnia and Herzegovina). The experiment was set up in a randomized block design with four pyrophyllite treatment rates i.e. 0, 200, 400 and 600 kg ha⁻¹ in three replications. Analyses of the heavy metals (Cu, Zn, Mn, Ni, Cr, Pb, Cd) in soil and plant samples were performed using atomic absorption spectrophotometry. Pyrophyllite addition in soil was found to reduce the availability of all tested heavy metals in the studied soil. The pyrophyllite addition at a rate of 200 kg ha⁻¹ reduced Mn, Cu and Zn available forms in soil by 11.1, 20.4 and 11.2%, respectively, compared with control. The pyrophyllite addition at higher rates i.e. 400 and 600 kg ha⁻¹ had an even higher impact on the decrease in Mn and Zn mobility in studied soil in comparison with 200 kg ha⁻¹. Additionally, these pyrophyllite rates have the ability to reduce Ni mobility in studied soil. The study also found a positive effect of all pyrophyllite treatments to reduce heavy metals accumulation in the leaves of potato grown on the studied soil. In sum, the results of this study indicate that pyrophyllite treatment could be an effective technique for improving the environmental quality of soils and alleviating the hazards of heavy metals to plants. However, further studies are necessary to confirm or denied this hypothesis.

Key words: accumulation, clay minerals, soil-plant system.

INTRODUCTION

Aluminosilicate clay minerals such as zeolite, pyrophyllite and bentonite have been widely studied regarding their suitability to be used in retention of heavy metal ions and thus for the protection, improvement and also remediation of soils polluted by heavy metals (Chaves et al., 2015; Lee et al., 2019). These minerals, based on a three-

dimensional aluminosilicate framework with numerous channels and cavities, have high retention properties which make them very useful in decreasing bioavailability of heavy metals in soils (Jemeljanova et al., 2019). Most scientists have noted that the ability of aluminosilicate minerals to reduce the mobility and thus availability of heavy metals in soils for plants is the result of their high cation exchange capacity, high surface area, and pore volume enabling the entry and retention of heavy metals in their inter layers (Wang & Li, 2011; Uddin, 2017; Xu et al., 2017).

The use of aluminosilicate minerals as heavy metals retention materials has advantages upon many other remediation techniques in terms of their non-toxic nature, low-cost, and high efficiency (Sharma et al., 2018). Among aluminosilicates, one of the least used materials in the soil remediation is pyrophyllite, which is a result of its relatively lower presence in nature, but also insufficient research on its use as a remediation material.

Pyrophyllite belongs to the class of aluminosilicates with form of the 2:1 layer i.e., the pyrophyllite structure consists of an octahedral Al-O layer sandwiched between two opposing tetrahedral Si-O layers. The bonding between these sandwiches is weak resulting with pyrophyllite's softness (Mohs hardness is between 1 and 1.5). Furthermore, the pyrophyllite has high density (2.7 and 2.9 g cm⁻³), a relatively high cation exchange capacity (between 50 and 70 meq 100 g⁻¹) and a pH ranging from neutral to slightly alkaline. It occurs in all shades of colour, particularly white, grey-white, and greenish pink, depending upon the presence of coloured minerals in pyrophyllite ore (Churakov, 2006).

Over the past several decades, the steel industry in Zenica has been associated with emission of large quantities of harmful substances, affecting negatively the air quality, as well as the quality of the soils in the Zenica area, more precisely their use value. Among harmful substances, heavy metals are of great concern due to its adverse effects on human health (Felix-Henningsen et al., 2010; Imeri et al., 2016). However, the presence of heavy metals in soils does not necessarily predict adverse effects on plants and consequently human health if they do not occur in a quantity above permissible limits established by legislation and in forms that is easily absorbed by plants (Kulokas et al., 2019). The normal range of Cu, Zn, Mn, Ni, Cr, Pb and Cd in leaves of plants are 5–20 mg kg⁻¹, 20–100 mg kg⁻¹, 15–150 mg kg⁻¹, 0.02–50 mg kg⁻¹, 0.1–1 mg kg⁻¹, 0.5–30 mg kg⁻¹ and 0.1–2.4 mg kg⁻¹, respectively (Chaney, 1989; Kastori et al., 1997).

The main objective of this study was to examine the ability of pyrophyllite ore materials from Parsovići – Konjic to decrease the availability of heavy metals in soils located near the steel mill in Zenica. An additional goal of this study was to evaluate pyrophyllite efficiency in reducing heavy metals accumulation in leaves of potato grown on these soils. Potato is food crops that are mostly grown in the study area and therefore is selected as the subject of this study.

MATERIALS AND METHODS

Materials

The pyrophyllite ore materials from the deposits 'Parsovići–Konjic' (Bosnia and Herzegovina) were used in this study. The particle size of the pyrophyllite materials used in the experiment was smaller than 500 µm.

The median total SiO₂, Al₂O₃, K, Ca, Mg, Cu, Ni, Zn, Co, Mn, Pb and Cr of pyrophyllite ore materials used in this study were 67.6%, 19.1%, 0.3%, 6.7%, 0.1%, 1.4 mg kg⁻¹, 2.7 mg kg⁻¹, 25.7 mg kg⁻¹, 0.4 mg kg⁻¹, 93.1 mg kg⁻¹, 8.0 mg kg⁻¹ and 0.8 mg kg⁻¹, respectively. The results mentioned above were obtained from the Laboratory at the Faculty of Agriculture and Food Sciences University of Sarajevo.

Study area

Three agricultural soil plots in Gradište, north-western suburb of Zenica (44°22'5" N, 17°89'85" E) were chosen as a study area (Fig. 1). The area has a temperate oceanic climate (cool summers and cool but not cold winters), with an average annual mean temperature and rainfall of 11.3 °C and 992 mm, respectively. It typically lacks a dry season as rainfall is dispersed evenly throughout the year.




Figure 1. Location map of the study site.

The investigated soil plots had approximately 1,000 m² in area and were located at a very close distance from each other (up to 500 m). These soils were chosen for investigation because it is located near the steel mill, but also because the populations in this area are fully or partially engaged in agriculture. Accordingly, the results of this study could provide specific data regarding the adverse impacts of heavy metals emission from steel factory on food crops production in the investigated area. All investigated soils were classified as Eutric Cambisol based on the World Reference Base for Soil Resources (FAO, 2014). Slightly acid to neutral reaction, medium texture, good physical properties, moderate organic matter content and base saturation of more than 50% is a typical characteristic of this type of soil.

Soil samples from investigated soil plots were collected once in March 2019 from 0–30 cm depth, using clean steel shovel. Five soil samples from each experimental plot were randomly gathered and mixed properly to obtain a composite soil sample. Thereafter, composite soil samples were placed in plastic bags and brought to the laboratory.

The soil samples were air-dried, crushed and ground using porcelain mortar and pestle, passed through a 2 mm and 1 mm sieve and then stored in paper bags until chemical analysis. Soil with highest concentrations of heavy metals (Cu, Zn, Mn, Ni, Cr, Pb, Cd) was used for conducting the experiment.

Experimental design

The experimental soil area was divided into twelve equal plots. Each plot had area 20 m² and the size of each unit plot was 5×4 m. The experiment was set up in a randomized block design with four pyrophyllite treatment in three replications. Distances between two plots were 1 m and the blocks were 2 m apart. Experiment treatments were as follows:

1. T₁ – soil without pyrophyllite i.e. control treatment
2. T₂ – soil with pyrophyllite at rate of 200 kg ha⁻¹
3. T₃ – soil with pyrophyllite at rate of 400 kg ha⁻¹
4. T₄ – soil with pyrophyllite at rate of 600 kg ha⁻¹.

Recommended pyrophyllite ore rate was recalculated based on experimental plot area (20 m²). Pyrophyllite ore materials in all experimental plots were applied seven days before planting potato. All other agrotechnical measures needed for optimum potato growth (pest control measures, irrigation) were performed identically on all experimental plots until the time of potato technological maturity.

Concentrations of plant-available forms of heavy metals in soils and heavy metal concentrations in potato leaves were determined at the end of experiment i.e. potato technological maturity stage. All samples of soils and potato leaves in the experiment area were collected at the same time.

Soil analysis

Before performing experiment, the following soil chemical properties were analysed: soil reaction (pH), organic matter (OM), available forms of phosphorus (available P) and potassium (available K), and total forms of heavy metals (Cu, Zn, Mn, Ni, Cr, Pb, Cd). Available forms of heavy metals in soil samples were determined at the end of experiment.

Soil pH was measured in H₂O and 1 M KCl in a 1:2.5 soil: solution ratio with a Mettler Toledo 320 pH meter. OM was measured by chromic acid digestion method (ISO 14235, 1998) and available forms of phosphorus and potassium by Egnér–Riehm method (Egnér et al., 1960).

Total heavy metals in soil samples were extracted by mineralizing 1 g of dry weight sample with 21 mL aqua regia (HNO₃: HCl 1:3) for 16 h at ambient temperature. Then, the flask solution was heated on hotplate under reflux for 2 h at 180 °C, cooled down to room temperature, filtered through quantitative filter paper into 100 mL flasks and diluted to the mark with deionized water (ISO 11466, 1995).

Plant-available forms of heavy metals in soil samples were extracted by EDTA solution (0.01 M ethylenediaminetetraacetic acid (EDTA) and 1 M (NH₄)₂CO₃, adjusted to pH 8.6) as follows: 10 g of air-dried soil was transferred into 100 mL plastic bottle, and then 20 mL EDTA solution was added. The bottle solution was shaken for 30 min at 180 rpm in an orbital shaker and thereafter the extract was filtered through quantitative filter paper into 25 mL flask and diluted to the mark with deionized water (Trierweiler & Lindsay, 1969).

Total and available heavy metal concentrations in soil samples were determined by atomic absorption spectrometry (ISO 11047, 1998) and their content expressed as mg per kg dry weight.

Plant sampling and analysis

Healthy green potato leaves without signs of parasites or disease from each experimental plot were carefully collected at the stage of potato technological maturity in quantity of approximately 200 g. The leaf samples were air-dried and separately powdered with a stainless-steel mill and stored in paper bags until analysis.

Total heavy metals in leaf samples were extracted by mineralizing 1 g of dry weight sample with 14 mL HNO₃ + HClO₄ mixture (2.5:1 v/v) for 4 h at ambient temperature. Thereafter, the flask solution was heated on a hotplate for 30 min, cooled down to room temperature, filtered through quantitative filter paper into 50 mL flasks and diluted to the mark with deionized water (Lisjak et al., 2009). Heavy metal concentrations in leaf samples were also determined by atomic absorption spectrometry (ISO 11047, 1998) and their content expressed as mg per g dry weight.

Statistical analysis

All measurements were done in triplicates and the results were presented as mean ± standard deviation. The collected data were analysed statistically using Microsoft Excel 2013 software program, and the significant differences between the variants were determined using *Least Significant Differences test* at 0.05 level of probability ($P \leq 0.05$).

RESULTS

Basic chemical properties of the studied soils

The results of the soil chemical analysis are presented in Table 1.

Chemical analyses showed that the sampled soils from studied were slight acidic to neutral with moderate levels of organic matter and high content of available forms of phosphorus (P₂O₅) and potassium (K₂O). Soil 1 had highest concentrations of heavy metals (Cu, Zn, Mn, Cr, Pb, Cd) and therefore this soil was selected for conducting the experiment.

Table 1. Chemical analysis of soil sample

Parameter	Unit	Soil 1	Soil 2	Soil 3
pH H ₂ O	pH unit	7.4	7.3	7.4
pH KCl	pH unit	6.7	6.6	6.8
organic matter	%	2.81	2.23	2.48
P ₂ O ₅	mg 100 g ⁻¹	20.84	8.08	3.22
K ₂ O	mg 100 g ⁻¹	60.9	30.3	34.9
Cu	mg kg ⁻¹	47.1	41.2	40.2
Zn	mg kg ⁻¹	53.9	56.7	50.0
Mn	mg kg ⁻¹	1,411.9	1,322.1	1,400.3
Ni	mg kg ⁻¹	120.2	160.0	155.9
Cr	mg kg ⁻¹	31.1	18.0	20.3
Pb	mg kg ⁻¹	69.5	54.1	55.0
Cd	mg kg ⁻¹	0.25	0.12	0.13

Available forms of heavy metals in the soil after pyrophyllite treatment

The concentrations of available forms of Cu, Zn, Mn, Pb, Ni, Cr and Cd in soil plots (mg kg⁻¹ dry weight), depending on the pyrophyllite treatment, are presented in Table 2 and 3. These results are also presented by a histogram 1 and 2 for easier visualization (Fig. 2).

The presented data illustrates the concentrations of available forms of heavy metals in soil after pyrophyllite treatment. The addition of pyrophyllite reduced the availability of all tested heavy metals in the studied soil compared to control treatments but the magnitude of the effect was not the same for all treatments. However, the statistical

analysis does not confirm the significant effect of added pyrophyllite on the reduction of Cr availability in soils.

Table 2. Concentrations of available forms of Cu, Zn, Mn and Pb in the studied soil

Treatment ¹	Cu	Zn	Mn	Pb
T ₁	3.83 ± 0.24 ^a	3.49 ± 0.15 ^a	7.87 ± 0.25 ^a	11.95 ± 1.61 ^a
T ₂	3.05 ± 0.50 ^b	3.10 ± 0.14 ^b	7.00 ± 0.26 ^b	4.20 ± 0.44 ^b
T ₃	2.89 ± 0.24 ^b	3.02 ± 0.08 ^b	5.89 ± 1.35 ^c	4.85 ± 1.42 ^b
T ₄	3.08 ± 0.36 ^b	3.01 ± 0.40 ^b	5.07 ± 0.78 ^d	5.73 ± 2.91 ^b
Lsd _{0.05}	0.451	0.203	0.713	1.851

Values marked with different letters in the same column indicate significantly differences. ¹Experimental treatment: T₁ – control treatment (without pyrophyllite); T₂ – pyrophyllite at rate of 200 kg ha⁻¹; T₃ – pyrophyllite at rate of 400 kg ha⁻¹; T₄ – pyrophyllite at rate of 600 kg ha⁻¹.

Table 3. Concentrations of available forms of Ni, Cr and Cd in the studied soil

Treatment ¹	Ni	Cr	Cd
T ₁	0.37 ± 0.03 ^a	0.107 ± 0.02	0.091 ± 0.02 ^a
T ₂	0.26 ± 0.04 ^b	0.106 ± 0.02	0.050 ± 0.01 ^b
T ₃	0.25 ± 0.05 ^b	0.105 ± 0.01	0.048 ± 0.01 ^b
T ₄	0.24 ± 0.03 ^b	0.105 ± 0.03	0.046 ± 0.01 ^b
Lsd _{0.05}	0.056	-	0.017

Values marked with different letters in the same column indicate significantly differences. ¹Experimental treatment: T₁ – control treatment (without pyrophyllite); T₂ – pyrophyllite at rate of 200 kg ha⁻¹; T₃ – pyrophyllite at rate of 400 kg ha⁻¹; T₄ – pyrophyllite at rate of 600 kg ha⁻¹.




Figure 2. The concentrations of available forms of heavy metals in soil depending on the pyrophyllite treatment.

Heavy metal concentrations in leaves of potato after pyrophyllite treatment

The heavy metal concentrations (Cu, Zn, Mn, Pb, Ni, Cr and Cd) in potato leaves (mg kg⁻¹ dry weight), depending on the pyrophyllite treatment, are presented in Table 4 and 5. These results are also presented by a histogram 3 and 4 for easier visualization (Fig. 3).

The presented data have shown that pyrophyllite treatment, regardless of the applied rates, significantly reduced accumulation of heavy metals (Cu, Zn, Mn, Pb, Ni and Cr) in potato leaves as compared to control treatment. The study also found a positive effect of pyrophyllite treatment to reduce Cd accumulation in potato leaves, but these findings did not reach statistical significance.

Table 4. Concentrations of Cu, Zn, Mn and Pb in potato leaves

Treatment ¹	Cu	Zn	Mn	Pb
T ₁	19.4 ± 5.6 ^a	44.5 ± 7.2 ^a	177.9 ± 69.2 ^a	18.0 ± 1.7 ^a
T ₂	12.5 ± 1.0 ^b	30.0 ± 4.6 ^b	78.3 ± 26.1 ^b	5.4 ± 2.9 ^b
T ₃	12.0 ± 2.9 ^b	28.1 ± 8.3 ^{bc}	71.5 ± 10.7 ^b	5.8 ± 1.8 ^b
T ₄	11.2 ± 0.7 ^b	24.6 ± 1.7 ^c	63.7 ± 11.1 ^b	3.8 ± 2.6 ^b
Lsd _{0.05}	2.94	5.24	35.34	2.14

Values marked with different letters in the same column indicate significant differences. ¹Experimental treatment: T₁ – control treatment (without pyrophyllite); T₂ – pyrophyllite at rate of 200 kg ha⁻¹; T₃ – pyrophyllite at rate of 400 kg ha⁻¹; T₄ – pyrophyllite at rate of 600 kg ha⁻¹.

Table 5. Concentrations of Ni, Cr and Cd in potato leaves

Treatment ¹	Ni	Cr	Cd
T ₁	10.6 ± 4.7 ^a	1.81 ± 0.32 ^a	0.24 ± 0.1
T ₂	3.0 ± 2.1 ^b	0.93 ± 0.36 ^b	0.24 ± 0.1
T ₃	2.5 ± 1.2 ^b	0.91 ± 0.61 ^b	0.23 ± 0.1
T ₄	2.1 ± 1.7 ^b	0.85 ± 0.24 ^b	0.20 ± 0.2
Lsd _{0.05}	2.50	0.35	–

Values marked with different letters in the same column indicate significant differences. ¹Experimental treatment: T₁ – control treatment (without pyrophyllite); T₂ – pyrophyllite at rate of 200 kg ha⁻¹; T₃ – pyrophyllite at rate of 400 kg ha⁻¹; T₄ – pyrophyllite at rate of 600 kg ha⁻¹.





Figure 3. Accumulation of heavy metals in soil depending on the pyrophyllite treatment.

DISCUSSION

This paper is an attempt to describe and evaluate the efficiency of pyrophyllite ore material to decrease the mobility of heavy metals in soil and thus their availability for plants. The results of this study demonstrate the high potential of pyrophyllite to immobilize heavy metals in studied soil, regardless of the amount of applied material. These findings are generally in line with previous studies (Kim et al., 2013; Park et al., 2017; Rath et al., 2017).

In the present study, the addition of pyrophyllite was especially successful in reducing Pb and Cd mobility in soil. The addition of pyrophyllite at rate of 600 kg ha⁻¹ reduced Pb and Cd available forms in soil by 52.1 and 49%, respectively, compared with control treatment. The effect of pyrophyllite at rate of 200 and 400 kg ha⁻¹ on the decrease Pb and Cd mobility in soil was also highly significant. These results are in agreement with results obtained by Singh et al. (2016).

Considering the soils polluted by Pb and Cd may pose hazards to food crops and consequently human health (Tchounwou et al., 2012; Alle et al., 2016; Borgulat et al.,

2018; Luo et al., 2019), the results mentioned above provide important scientific information in an effort to improve soil protection and quality by applying pyrophyllite. Our opinion is pyrophyllite has the potential to be included in the inventory of efficient soil remediation techniques, especially because the use of pyrophyllite for soil remediation is simple to operate, cost effective and very reliable.

Osacky et al. (2015) reported that the ability of pyrophyllite and other clay minerals to reduce the mobility of heavy metals in soils is the result of their potential to form complexes with heavy metals in their inter layers or on surface areas. The pyrophyllite primarily binds heavy metals in the space between the layers, due to their large pore volume (Ismadji et al., 2015). The efficiency of pyrophyllite to immobilize heavy metals in soil also depends on their dosage, surface area as well as soil physical and chemical properties.

Furthermore, the speciation or chemical form of the element play an important role in in the fixation of heavy metals on the pyrophyllite surface. Namely, heavy metals often have different levels of mobility depending on the specific metal oxidation state (Violante et al., 2010). For example, Cr (III) is, in general, much less toxic and mobile in soils than Cr (VI), and therefore reactions that reduce Cr (VI) to Cr (III) are of great importance for soil remediation.

Although numerous studies confirm that pyrophyllite has high efficiency in removing heavy metals from the soil and water (Prasad & Saxena, 2008; Chawla et al., 2018; Panda et al., 2018), the heavy metal binding mechanisms on pyrophyllite are not completely clear. Scheidegger et al. (1996) have attempted to explain the binding mechanisms between Ni and pyrophyllite surfaces. They reported pH is the primary factor that controls Ni binding on pyrophyllite. The study found that Ni sorption on pyrophyllite, in the lower pH region (i.e., $\text{pH} < 7$), increased with decreasing ionic strength. Contrary, Ni sorption on pyrophyllite, in the higher pH region ($\text{pH} > 7$ with high Ca^{2+} level), was slower. These results can be explained by the fact that Ni with other ions (primarily Ca^{2+}) compete for the same free sites on the pyrophyllite adsorptive complex. Similar results were reported by Gou et al. (2018).

The results of this study also showed that pyrophyllite addition at a rate of 400 and 600 kg ha^{-1} in soil has the ability to reduce Ni mobility in studied soil (neutral to weakly basic), indicating the pyrophyllite possesses some mechanisms for Ni retention even in the higher pH region. Zhao et al. (2017) noted the inner-sphere complexes between Ni and pyrophyllite surface areas (with no intervening water molecules) were dominant mechanisms for Ni retention.

In the present study, pyrophyllite also demonstrated the high potential to immobilize Mn, Cu and Zn in studied soil. Namely, the pyrophyllite addition at a rate of 200 kg ha^{-1} reduced Mn, Cu and Zn available forms in soil by 11.1, 20.4 and 11.2%, respectively, compared with control treatment (without pyrophyllite). The pyrophyllite addition at higher rates i.e. 400 and 600 kg ha^{-1} had an even higher impact on the decrease in Mn and Zn mobility in studied soil in comparison with 200 kg ha^{-1} pyrophyllite rate. These findings indicate that pyrophyllite ore material used in this research is potentially useful additive to bind Mn and Zn ions in soils and that their mobility in soil decreases with increasing pyrophyllite rates.

However, a better understanding of the pyrophyllite sorption mechanism could make a significant contribution to improving pyrophyllite efficiency to reduce heavy metals mobility in soils. An interesting data related to pyrophyllite is the fact that

pyrophyllite has the ability to easily disperse in water, enabling a higher area of pyrophyllite exposure in the soil and thus its activity (El Gaidoumi et al., 2019), and finally pyrophyllite could increase soil pH, thus resulting in less heavy metal mobility (Newton & Sposito, 2015). All the above-mentioned scientific data are undoubtedly associated with its sorption mechanism.

As shown in Table 3, all pyrophyllite treatments were also significantly reduced the accumulation of heavy metals (Cu, Zn, Mn, Ni, Cr, Pb and Cr) in the leaves of potato grown on the studied soil. In addition, the pyrophyllite treatments reduced Cd accumulation in potato leaves, but these effects were not statistically significant. The results also showed that 600 kg ha⁻¹ pyrophyllite rate had the highest effect on the reduction of heavy metals accumulation in potato leaves. This treatment reduced the accumulation of Cu, Zn, Mn, Ni, Cr, Pb and Cr in the potato leaves by 42.1, 44.5, 64.2, 80.0, 52.8, 78.9 and 17.8%, respectively, compared to control treatment. The effect of 200 and 400 kg ha⁻¹ pyrophyllite rate was also significant but less pronounced.

Generally, the results of this study demonstrated that the addition of pyrophyllite in tested soil reduced heavy metals mobility, thus resulting with low accumulation of heavy metals in potato leaves. One exception to this general rule was the behaviour of Cr in soil- plant system. Namely, the pyrophyllite treatments did not significantly reduce the concentration of Cr available forms in studied soil at the end of experiment, but significantly reduced its accumulation in the leaves of potato grown in the same soil plots. This inconsistent result could be potentially attributed to Cr speciation. As is well known, the Cr mobility in soil can change drastically depending on change in soil pH, redox potential, microbial activity, content of organic matter, content and type of clay minerals etc. Accordingly, changes in Cr chemical forms (speciation) are possible in a relatively short-term, and thus its bioavailability (Huang et al., 2018). We assume that Cr mobility in soil in the initial stages of potato growth were higher, resulting in higher Cr uptake by the potato. However, further studies are necessary to confirm or denied this hypothesis as well as other hypotheses presented in this study.

CONCLUSIONS

Pyrophyllite addition in soil was found to reduce the availability of tested heavy metals in the studied soil, indicating that pyrophyllite treatment could be an effective technique to stabilize soils polluted by heavy metals. Under experimental conditions of the study, the pyrophyllite addition at a rate of 200 kg ha⁻¹ reduced Mn, Cu and Zn available forms in soil by 11.1, 20.4 and 11.2%, respectively, compared with control. The pyrophyllite addition at higher rates i.e. 400 and 600 kg ha⁻¹ had an even higher impact on the decrease in Mn and Zn mobility in studied soil in comparison with 200 kg ha⁻¹. Additionally, the accumulation of all tested heavy metals in leaves of potato grown on soil plots treated with pyrophyllite was found to be lower than those in non-treated plots, indicating that pyrophyllite can also alleviate the risk for plants associated with heavy metals.

ACKNOWLEDGEMENTS. This study was financially supported by the Environmental Fund of the Federation of Bosnia and Herzegovina.

REFERENCES

- Alle, V., Kondratovics, U., Osvalde, A. & Vikmane, M. 2016. Differences in cadmium accumulation and induced changes in root anatomical structures in plants used for food. *Agronomy Research* **14**(S2), 1249–1260.
- Borgulat, J., Mętrak, M., Staszewski, T., Wilkomirski, B. & Suska-Malawska, M. 2018. Heavy Metals Accumulation in Soil and Plants of Polish Peat Bogs. *Polish Journal of Environmental Studies* **27**(2), 537–544.
- Chaney, R.L. 1989. Toxic element accumulation in soils and crops: Protecting soil fertility and agricultural foodchains. In Bar-Yosef, B., Barrow, N.J. & Goldshmid, J. (ed.): *Inorganic contaminants in the Vadose Zone*. Springer, Berlin, pp. 140–158.
- Chaves, L.H.G, Tito, G.A. & Chaves, B. 2015. Bentonite Application in the Remediation of Zinc Contamination Soil. *Chemical Engineering Transactions* **58**, 745–750.
- Chawla, A., Prasad, M., Goswami, R., Ranshore, S., Kulshreshtha, A. & Sinha, A.S.K. 2016. Kinetic model for sorption of divalent heavy metal ions on low cost minerals. *Korean Journal of Chemical Engineering* **33**(2), 649–656.
- Churakov, S.V. 2006. Ab Initio Study of Sorption on Pyrophyllite: Structure and Acidity of the Edge Sites. *The Journal of Physical Chemistry B* **110**(9), 4135–4146.
- Egnér, H., Riehm, H. & Domingo, W.R. 1960. Studies on the chemical soil analysis as a basis for the assessment of the nutrient status of soils. II. Chemical extraction methods for determination of phosphorus and potassium. *Kunliga Lantbrukshögskolans Annaler* **26**, 199–215 (in German).
- El Gaidoumi, A., Doña-Rodríguez, J.M. & Melián, E.P. 2019. Mesoporous pyrophyllite-titania nanocomposites: synthesis and activity in phenol photocatalytic degradation. *Research on Chemical Intermediates* **45**(2), 333–353.
- Felix-Henningsen, P., Urushadze, T., Steffens, D., Kalandadze, B. & Narimanidze, E. 2010. Uptake of heavy metals by food crops from highly-polluted Chernozem-like soils in an irrigation district south of Tbilisi, eastern Georgia. *Agronomy Research* **8**(1), 781–795.
- FAO. 2014. 'International soil classification system for naming soils and creating legends for soil maps.' Word Soils Resources Reports, Rome, Italy, 2014. Available at: <http://www.fao.org/3/i3794en/i3794en.pdf>
- Gou, W., Siebecker, M.G., Wang, Z. & Li, W. 2018. Competitive sorption of Ni and Zn at the aluminum oxide/water interface: an XAFS study. *Geochemical Transactions* **19**(1), 9.
- Huang, W., Jiao, J., Ru, M., Bai, Z., Yuan, H., Bao, Z. & Liang, Z. 2018. Localization and Speciation of Chromium in *Coptis chinensis* Franch. using Synchrotron Radiation X-ray Technology and Laser Ablation ICP-MS. *Scientific Reports* **8**(1), 8603.
- Imeri, R., Kullaj, E., Duhani, E. & Millaku, L. 2016. Impact of rootstock on heavy metal bioaccumulation in apple plant grown near an industrial source in Obiliq, Kosovo. *Agronomy research* **17**(1), 100–110.
- Ismadji, S., Soetaredjo, F.E. & Ayucitra, A. 2015. *Clay Materials for Environmental Remediation*. SpringerBriefs in Green Chemistry for Sustainability. Springer, UK, 113 pp.
- ISO 11047. 1998. 'Determination of cadmium, chromium, cobalt, copper, lead, manganese, nickel and zinc - Flame and electrothermal atomic absorption spectrometric methods.' International Organization for Standardization, Geneva, Switzerland. Available at: <https://www.iso.org/standard/24010.html>
- ISO 11466. 1995. 'Extraction of trace elements soluble in aqua regia.' International Organization for Standardization, Geneva, Switzerland. Available at: <https://www.iso.org/standard/19418.html>

- ISO 14235. 1998. 'Soil quality – Determination of organic carbon in soil by sulfochromic oxidation.' International Organization for Standardization, Geneva, Switzerland. Available at: <https://www.iso.org/standard/23140.html>
- Jemeljanova, M., Ozola, R. & Klavins, M. 2019. Physical-chemical properties and possible applications of clay minerals and humic acid composite materials. *Agronomy research* **17**(S1), 1023–1032. doi.org/10.15159/AR.19.019
- Kastori, R., Petrović, N. & Arsenijević-Maksimović, I. 1997. Heavy metals and plants. In: Kastori, R. (ed): *Heavy Metals in the Environment*, pp. 195–257 (in Serbian).
- Kim, J., Lee, C., Park, J., Kang, J., Choi, N. & Kim, S. 2013. Use of pyrophyllite clay for fluoride removal from aqueous solution. *Desalination and Water Treatment* **51**(16), 3408–3416.
- Kulokas, M., Zaleskas, V., Pedišius, N., Praspaliauskas, M. & Buinevičius, K. 2019. Properties of biofuel fly ash and capabilities of its use for agricultural needs. *Agronomy research* **17**(2), 540–550. doi.org/10.15159/ar.19.051
- Lee, D.S., Lim, S.S., Park, H.J., Yang, H.I., Park, S.I., Kwak, J.H. & Choi, W.J. 2019. Fly ash and zeolite decrease metal uptake but do not improve rice growth in paddy soils contaminated with Cu and Zn. *Environment International* **129**, 551–564.
- Lisjak, M., Spoljarevic, M., Agic, D. & Andric, L. 2009. *Practical Plant Physiology*. Faculty of Agriculture, Osijek, 25 pp. (in Croatian).
- Luo, K., Liu, H., Zhao, Z., Long, J., Li, J., Jiang, C. & Rao, C. 2019. Spatial Distribution and Migration of Cadmium in Contaminated Soils Associated with a Geochemical Anomaly: A Case Study in Southwestern China. *Polish Journal of Environmental Studies* **28**(5), 3799–3807.
- Newton, A.G. & Sposito, G. 2015. Molecular Dynamics Simulations of Pyrophyllite Edge Surfaces: Structure, Surface Energies, and Solvent Accessibility. *Clays and Clay Minerals* **63**(4), 277–289.
- Osacky, M., Geramian, M., Ivey, D.G., Liu, Q. & Etsell, T.H. 2015. Influence of Nonswelling Clay Minerals (Illite, Kaolinite, and Chlorite) on Nonaqueous Solvent Extraction of Bitumen. *Energy Fuels* **29**(7), 4150–4159.
- Panda, L., Rath, S.S., Rao, D.S., Nayak, B.B., Das, B. & Misra, P.K. 2018. Thorough understanding of the kinetics and mechanism of heavy metal adsorption onto a pyrophyllite mine waste based geopolymer. *Journal of Molecular Liquids* **263**, 428–441.
- Park, J., Kang, J. & Kim, S. 2017. Comparative Analysis of Bacteriophages and Bacteria Removal in Soils and Pyrophyllite-Amended Soils: Column Experiments. *Water, Air & Soil Pollution* **228**(3), 103.
- Prasad, M. & Saxena, S. 2008. Attenuation of divalent toxic metal ions using natural sericitic pyrophyllite. *Journal of Environmental Management* **88**(4), 1273–1279.
- Rath, S.S., Singh, S., Rao, D.S., Nayak, B.B. & Mishra, B.K. 2017. Adsorption of heavy metals on a complex Al-Si-O bearing mineral system: Insights from theory and experiments. *Separation and Purification Technology* **186**, 28–38.
- Scheidegger, A.M., Sparks, D.L. & Fendorf, M. 1996. Mechanisms of Nickel Sorption on Pyrophyllite: Macroscopic and Microscopic Approaches. *Soil Science Society of America Journal* **60**(6), 1763–1772.
- Sharma, S., Tiwari, S., Hasan, A., Saxena, V. & Pandey, L.M. 2018. Recent advances in conventional and contemporary methods for remediation of heavy metal-contaminated soils. *3 Biotech.* **8**(4), 216.
- Singh, S., Jena, S.K. & Das, B. 2016. Application of pyrophyllite mine waste for the removal of cadmium and lead ions from aqueous solutions. *Desalination and Water Treatment* **57**(19), 8952–8966.
- Tchounwou, P.B., Yedjou, C.G., Patlolla, A.K. & Sutton, D.J. 2012. Heavy metal toxicity and the environment. *EXS* **101**, 133–164.

- Trierweiler, J.E. & Lindsay, W.L. 1969. EDTA-ammonium carbonate soil test for zinc. *Soil Science Society of America Proceedings* **39**, 49–54.
- Uddin, M.K. 2017. A review on the adsorption of heavy metals by clay minerals, with special focus on the past decade. *Chemical Engineering Journal* **308**, 438–462.
- Violante, A., Cozzolino, V., Perelomov, L., Caporale, A.G. & Pigna, M. 2010. Mobility and bioavailability of heavy metals and metalloids in soil environments. *Journal of Soil Science and Plant Nutrition* **10**(3), 268–292.
- Wang, X.L. & Li, Y. 2011. Measurement of Cu and Zn adsorption onto surficial sediment components: New evidence for less importance of clay minerals. *Journal of Hazardous Materials* **189**(3), 719.
- Xu, Y., Liang, X., Xu, Y., Qin, X., Huang, Q., Wang, L. & Sun, Y. 2017. Remediation of heavy metal-polluted agricultural soils using clay minerals: A review. *Pedosphere* **27**(2), 193–204.
- Zhao, X., Qiang, S., Wu, H., Yang, Y., Shao, D., Fang, L., Liang, J., Li, P. & Fan, Q. 2017. Exploring the Sorption Mechanism of Ni (II) on Illite: Batch Sorption, Modelling, EXAFS and Extraction Investigations. *Scientific Reports* **7**, 8495.